arXiv:hep-th/9208037v1 13 Aug 1992

CERN–TH./92
DFTT-44/92

BICOVARIANT DIFFERENTIAL GEOMETRY OF THE QUANTUM GROUP $GL_q(3)$

Paolo Aschieri$^{\diamond *}$ and Leonardo Castellani*

$^{\diamond}$ *CERN, CH-1211 Geneva 23, Switzerland.*
**Istituto Nazionale di Fisica Nucleare, Sezione di Torino*
and
Dipartimento di Fisica Teorica
Via P. Giuria 1, 10125 Torino, Italy.

Abstract

We construct a bicovariant differential calculus on the quantum group $GL_q(3)$, and discuss its restriction to $[SU(3) \otimes U(1)]_q$. The q-algebra of Lie derivatives is found, as well as the Cartan-Maurer equations. All the quantities characterizing the non-commutative geometry of $GL_q(3)$ are given explicitly.

CERN–TH./92
DFTT-44/92
July1992

---

email addresses: Decnet=(39163::ASCHIERI, CASTELLANI; VXCERN::ASCHIERI)
Bitnet=(ASCHIERI, CASTELLANI@TORINO.INFN.IT)

It is tempting to use quantum groups [1]–[4] for the construction of generalized gauge and gravity theories. Some initial work in this direction can be found in refs. [5]-[8] for q-gauge theories and in [9], [21] for q-gravity theories. The basic idea of ref. [9] is to obtain a q-gravity theory as a "gauge" theory of the quantum Poincaré group (or of the quantum De-Sitter group). For this we need to study the differential (non-commutative) geometry of the relevant q-Poincaré (or q-De Sitter) group. As a first step, the bicovariant calculus on the $D=2$ q-Poincaré group was developed in [9].

On the other hand, the differential geometry of the quantum $SU(2)$ and $SU(3)$ groups is of interest in the construction of q-deformed gauge theories that reduce to the standard model for $q \to 1$. The Cartan-Maurer equations for the q-analogue of the left-invariant one-forms *define* the field strengths, as in the classical case, and one can find an action of the type $F^i_{\mu\nu} F^j_{\mu\nu} g_{ij}$ invariant under the q-group symmetries [7].

The differential calculus on q-groups, initiated by Woronowicz [10], has been developed in a number of papers from various points of view [11]-[22]. The general theory being still in its infancy, we feel that explicit non-trivial examples may be inspiring, and in this Letter we present a (bicovariant) differential calculus on $GL_q(3)$ and on its restriction to $[SU(3) \otimes U(1)]_q$.

The quantum group $GL_q(3)$ is defined [3] as the (associative) algebra A freely generated by:

i) non-commuting matrix entries ${T^a}_b$, a,b=1,2,3, satisfying

$$R^{ab}{}_{ef} T^e{}_c T^f{}_d = T^b{}_f T^a{}_e R^{ef}{}_{cd} \tag{1}$$

where the R matrix is given in Table 1, the rows and columns being numbered in the order $11, 12, 13, 21, 22, 23, 31, 32, 33$. The R matrix satisfies the quantum Yang-Baxter equation

$$R^{a_1 b_1}{}_{a_2 b_2} R^{a_2 c_1}{}_{a_3 c_2} R^{b_2 c_2}{}_{b_3 c_3} = R^{b_1 c_1}{}_{b_2 c_2} R^{a_1 c_2}{}_{a_2 c_3} R^{a_2 b_2}{}_{a_3 b_3}, \tag{2}$$

a sufficient condition for the consistency of the "RTT" relations (1). One can also think of the indices ${}^a{}_b$ of ${T^a}_b$ as composite indices taking the values $1, ...9$ (with $11 \to 1,\ 12 \to 2,\ 13 \to 3,\ 21 \to 4,\ 22 \to 5,\ 23 \to 6,\ 31 \to 7,\ 32 \to 8,\ 33 \to 9$), so that

$$T^a{}_b = \begin{pmatrix} T_1 & T_2 & T_3 \\ T_4 & T_5 & T_6 \\ T_7 & T_8 & T_9 \end{pmatrix} \tag{3}$$

Then the relations (1) take the explicit form:

$$\begin{array}{lll}
T_1T_2 = qT_2T_1, & T_1T_3 = qT_3T_1, & T_2T_3 = qT_3T_2, \\
T_1T_4 = qT_4T_1, & T_1T_5 = (q-q^{-1})T_4T_2 + T_5T_1, & T_1T_6 = (q-q^{-1})T_4T_3 + T_6T_1, \\
T_2T_5 = qT_5T_2, & T_2T_6 = (q-q^{-1})T_5T_3 + T_6T_2, & T_1T_9 = (q-q^{-1})T_7T_3 + T_9T_1, \\
T_2T_8 = qT_8T_2, & T_2T_9 = (q-q^{-1})T_8T_3 + T_9T_2, & T_3T_9 = qT_9T_3, \\
T_4T_5 = qT_5T_4, & T_4T_6 = qT_6T_4, & T_5T_6 = qT_6T_5, \\
T_4T_7 = qT_7T_4, & T_4T_8 = (q-q^{-1})T_7T_5 + T_8T_4, & T_4T_9 = (q-q^{-1})T_7T_6 + T_9T_4, \\
T_5T_8 = qT_8T_5, & T_5T_9 = (q-q^{-1})T_8T_6 + T_9T_5, & T_6T_9 = qT_9T_6, \\
T_7T_8 = qT_8T_7, & T_7T_9 = qT_9T_7, & T_8T_9 = qT_9T_8.
\end{array} \tag{4}$$

all other commutations being trivial. For $q \rightarrow 1$ all elements commute (the $R^{ab}{}_{cd}$ matrix becomes $\delta^a_c \delta^b_d$, cf. Table 1).

ii) the identity element I and the inverse ξ of the q-determinant of T, defined by:

$$\xi \det_q T = \det_q T \xi = I \tag{5}$$

$$\det_q T \equiv \sum_\sigma (-q)^{l(\sigma)} T^1{}_{\sigma(1)} \cdots T^n{}_{\sigma(n)} \tag{6}$$

where $l(\sigma)$ is the minimum number of transpositions in the permutation σ. Explicitly:

$$\det_q T = T_1T_5T_9 + q^2T_2T_6T_7 + q^2T_3T_4T_8 - qT_1T_6T_8 - q^3T_3T_5T_7 - qT_2T_4T_9\ . \tag{7}$$

It is not difficult to check that ξ and $\det_q T$ commute with all the elements $T^a{}_b$. The non-commutative algebra generated by $T^a{}_b$, I and ξ is a Hopf algebra, i.e. we can define a coproduct Δ, a counit ε and a coinverse κ:

$$\Delta(T^a{}_b) = T^a{}_c \otimes T^c{}_b, \quad \Delta(I) = I \otimes I \tag{8}$$

$$\varepsilon(T^a{}_b) = \delta^a_b, \quad \varepsilon(I) = 1 \tag{9}$$

$$\kappa(T^a{}_b) = (T^{-1})^a{}_b, \quad \kappa(I) = I. \tag{10}$$

the (matrix) inverse of T being given by(cf. [3]):

$$(T^{-1})^a{}_b = \xi \left(\begin{array}{ccc}
T_5T_9 - qT_6T_8 & -q^{-1}T_2T_9 + T_3T_8 & q^{-2}T_2T_6 - q^{-1}T_3T_5 \\
-qT_4T_9 + q^2T_6T_7 & T_1T_9 - qT_3T_7 & -q^{-1}T_1T_6 + T_3T_4 \\
q^2T_4T_8 - q^3T_5T_7 & -qT_1T_8 + q^2T_2T_7 & T_1T_5 - qT_2T_4
\end{array} \right) \tag{11}$$

From eqs. (7) and (5) one deduces the co-structures on $\det_q T$ and ξ:

$$\Delta(\det_q T) = \det_q T \otimes \det_q T, \quad \Delta(\xi) = \xi \otimes \xi \tag{12}$$

$$\varepsilon(\det_q T) = 1, \quad \varepsilon(\xi) = 1 \tag{13}$$

$$\kappa(\det_q T) = \xi, \quad \kappa(\xi) = \det_q T \tag{14}$$

For $q \in R$ a q-analogue of complex conjugation ($*$-structure) can be consistently defined on A as [3]:

$$T^* = (T^{-1})^t \tag{15}$$

This unitarity condition restricts $GL_q(3)$ to its quantum subgroup $[SU(3) \otimes U(1)]_q$. The further constraint $\det_q T = I$ yields $SU_q(3)$. Other real forms of $GL_q(3)$ can be considered, via a more general unitarity condition [3].

The differential calculus we are about to construct can be entirely formulated in terms of the R matrix. This is true for all quantum groups of the A, B, C, D series. The general constructive procedure can be found in ref. [17], or, in the notations we adopt here, in ref. [21]. There the reader can find the derivation of the general formulas appearing later in this Letter.

As discussed in [10] and [17], we can start by introducing the (quantum) left-invariant one-forms ${\omega_a}^b$, whose exterior product

$${\omega_{a_1}}^{a_2} \wedge {\omega_{d_1}}^{d_2} \equiv {\omega_{a_1}}^{a_2} \otimes {\omega_{d_1}}^{d_2} - \Lambda_{a_1}{}^{a_2}{}_{d_1}{}^{d_2}|^{c_1}{}_{c_2}{}^{b_1}{}_{b_2} {\omega_{c_1}}^{c_2} \otimes {\omega_{b_1}}^{b_2} \tag{16}$$

is defined by the braiding matrix Λ:

$$\Lambda_{a_1}{}^{a_2}{}_{d_1}{}^{d_2}|^{c_1}{}_{c_2}{}^{b_1}{}_{b_2} \equiv d^{f_2} d^{-1}_{c_2} R^{f_2 b_1}{}_{c_2 g_1} (R^{-1})^{c_1 g_1}{}_{e_1 a_1} (R^{-1})^{a_2 e_1}{}_{g_2 d_1} R^{g_2 d_2}{}_{b_2 f_2} \tag{17}$$

For $q \rightarrow 1$ the braiding matrix Λ becomes the usual permutation operator and one recovers the classical exterior product.

In the case of $GL_q(3)$ there are nine independent left-invariant one-forms ${\omega_a}^b$, a,b=1,2,3, and we number them ω^i, i=1,...9, according to the same index convention used for the matrix elements $T^a{}_b$, i.e. ${\omega_1}^1 \rightarrow \omega^1, {\omega_1}^2 \rightarrow \omega^2$ etc. The d vector in (17) is given by $d_1 = q, d_2 = q^3, d_3 = q^5$.

The commutation relations for exterior products of ${\omega_a}^b$ are deduced from the formula [valid for all $GL_q(N)$], cf. [21]:

$$\omega^i \wedge \omega^j = -Z^{ij}{}_{kl} \omega^k \wedge \omega^l \tag{18}$$

$$Z^{ij}{}_{kl} \equiv \frac{1}{q^2 + q^{-2}} [\Lambda^{ij}{}_{kl} + (\Lambda^{-1})^{ij}{}_{kl}]. \tag{19}$$

For $GL_q(3)$, these relations are collected in Table 2 (we have omitted the wedge symbol in the exterior products between the ω).

As in the classical case, the left-invariant 1-forms ${\omega_a}^b$ are a basis for the space Γ of quantum 1-forms, i.e. any $\rho \in \Gamma$ can be expressed as $\rho = {a_b}^a {\omega_a}^b$, with ${a_b}^a \in A$.

The left and right action of the quantum group on Γ are defined as follows:

$$\Delta_L({\omega_{a_1}}^{a_2}) = I \otimes {\omega_{a_1}}^{a_2}, \tag{20}$$

$$\Delta_R({\omega_{a_1}}^{a_2}) = {\omega_{b_1}}^{b_2} \otimes M^{b_1}{}_{b_2 a_1}{}^{a_2}, \tag{21}$$

where $M^{b_1}_{b_2a_1}{}^{a_2}$, the adjoint representation, is given by

$$M^{b_1}_{b_2a_1}{}^{a_2} \equiv T^{b_1}{}_{a_1} \kappa(T^{a_2}{}_{b_2}). \tag{22}$$

The q-analogue of the fact that left and right actions commute is given by the property

$$(id \otimes \Delta_R)\Delta_L = (\Delta_L \otimes id)\Delta_R \tag{23}$$

The bimodule [1] Γ is further characterized by the commutations between $\omega_a{}^b$ and the basic elements of A:

$$\omega_{a_1}{}^{a_2} T^{b_1}{}_{b_2} = s\, T^{b_1}{}_d (R^{-1})^{dc_1}{}_{ea_1} (R^{-1})^{a_2e}{}_{c_2b_2} \omega_{c_1}{}^{c_2} \tag{24}$$

$$\omega_{a_1}{}^{a_2} \mathrm{det}_q T = s^3 q^{-2}\ \mathrm{det}_q T \omega_{a_1}{}^{a_2} \quad , \qquad \omega_{a_1}{}^{a_2} \xi = s^{-3} q^2\ \xi \omega_{a_1}{}^{a_2} \tag{25}$$

where s is a complex arbitrary scale (whose classical limit is 1), cf. refs. [20] and [21]. From eq. (25) we see that $s = q^{\frac{2}{3}}$ is a "canonical" choice for s (then $\mathrm{det}_q T$ and its inverse ξ commute with any element of Γ). Equations (24) also allow to find the commutations between a generic $\rho \in \Gamma$ and a generic $a \in A$, since any ρ is a linear combination of $\omega_a{}^b$ with coefficients belonging to A, and any a is a polynomial in the $T^a{}_b$.

Table 3 contains a list of the commutations (24) for $GL_q(3)$, with $s = 1$. The case $s \neq 1$ is recovered by multiplying the right-hand sides of Table 3 by s.

The induced $*$-structure on the ω's is given by [16], [17]:

$$(\omega_{a_1}{}^{a_2})^* = -\omega_{a_2}{}^{a_1} \tag{26}$$

and is extended to Γ via the rules

$$(a\rho)^* = \rho^* a^*, \quad (\rho a)^* = a^* \rho^*. \tag{27}$$

This $*$-structure is compatible with the left and right action of the quantum group, i.e.

$$\Delta(a^*) = [\Delta(a)]^*, \quad \forall a \in A \tag{28}$$

$$\Delta_L(\rho^*) = [\Delta_L(\rho)]^*, \quad \Delta_R(\rho^*) = [\Delta_R(\rho)]^*, \quad \forall \rho \in \Gamma \tag{29}$$

where $(a \otimes b)^* \equiv b^* \otimes a^*,\ (a \otimes \rho)^* \equiv a^* \otimes \rho^*,\ (\rho \otimes a)^* \equiv \rho^* \otimes a^*, \quad \forall a, b \in A,\ \rho \in \Gamma$.

The exterior differential on $\Gamma^{\wedge k} = \Gamma \wedge \Gamma \wedge \cdots \wedge \Gamma$ (k-times) is defined by means of the bi-invariant (i.e. left- and right-invariant) element $\tau = \sum_a \omega_a{}^a \in \Gamma$ as follows:

$$d\theta \equiv \frac{1}{q - q^{-1}} [\tau \wedge \theta - (-1)^k \theta \wedge \tau], \tag{30}$$

---

[1] "bimodule" meaning that $\rho \in \Gamma$ can be multiplied on the right and on the left by elements $a \in A$. In general $a\rho \neq \rho a$.

where $\theta \in \Gamma^{\wedge k}$. The normalization $\frac{1}{q-q^{-1}}$ is necessary in order to obtain the correct classical limit. For $a \in A$ we have

$$da = \frac{1}{q-q^{-1}}[\tau a - a\tau]. \tag{31}$$

This linear map satisfies the Leibniz rule

$$d(ab) = (da)b + a(db), \quad \forall a, b \in A; \tag{32}$$

and has the properties

$$d(\theta \wedge \theta') = d\theta \wedge \theta' + (-1)^k \theta \wedge d\theta' \tag{33}$$

$$d(d\theta) = 0 \tag{34}$$

$$d(\theta^*) = (d\theta)^* \tag{35}$$

$$\Delta_L(d\theta) = (id \otimes d)\Delta_L(\theta) \tag{36}$$

$$\Delta_R(d\theta) = (d \otimes id)\Delta_R(\theta), \tag{37}$$

where $\theta \in \Gamma^{\wedge k}$, $\theta' \in \Gamma^{\wedge k'}$. The $*$-structure on $\Gamma^{\wedge n}$ can be inferred from the rule

$$(\theta \wedge \theta')^* = (-1)^{kk'} \; \theta'^* \wedge \theta^* \tag{38}$$

The properties (36), (37) express the fact that d commutes with the left and right action of the quantum group, as in the classical case. Moreover, for a generic $\rho \in \Gamma$ we have

$$\rho = a_k db_k, \qquad\qquad a_k, b_k \in A \tag{39}$$

The relations of Table 6 can be inverted to yield the ${\omega_a}^b$ in terms of the ${dT^c}_d$, thus proving the decomposition (39).

The bimodule Γ and the mappings d, Δ_R, Δ_L with the above properties define a *bicovariant* differential calculus on the q-group [10].

The "quantum Lie algebra generators" are linear functionals on A (i.e. belonging to its dual A') introduced via the formula [10]

$$da = \frac{1}{q-q^{-1}}[\tau a - a\tau] = (\chi^{a_1}{}_{a_2} * a)\omega_{a_1}{}^{a_2}. \tag{40}$$

where

$$\chi * a \equiv (id \otimes \chi)\Delta(a), \qquad \forall a \in A, \; \chi \in A' \tag{41}$$

By taking the exterior derivative of (40), requiring that $d^2 = 0$ and using the fact that $\tau = {\omega_b}^b$ is bi-invariant, we arrive at the q-Lie algebra relations [17], [21]:

$$\chi^{d_1}{}_{d_2}\chi^{c_1}{}_{c_2} - \Lambda_{e_1}{}^{e_2}{}_{f_1}{}^{f_2}|^{d_1}{}_{d_2}{}^{c_1}{}_{c_2} \; \chi^{e_1}{}_{e_2}\chi^{f_1}{}_{f_2} = \mathbf{C}^{d_1}{}_{d_2}{}^{c_1}{}_{c_2}|_{a_1}{}^{a_2}\chi^{a_1}{}_{a_2} \tag{42}$$

where the structure constants are explicitly given by:

$$\mathbf{C}^{a_1}{}_{a_2}{}^{b_1}{}_{b_2}|_{c_1}^{c_2} = \frac{1}{q-q^{-1}}[-\delta^{b_1}_{b_2}\delta^{a_1}_{c_1}\delta^{c_2}_{a_2} + \Lambda_b{}^b{}_{c_1}{}^{c_2}|^{a_1}{}_{a_2}{}^{b_1}{}_{b_2}]. \tag{43}$$

and $\chi^{d_1}{}_{d_2}\chi^{c_1}{}_{c_2} \equiv (\chi^{d_1}{}_{d_2} \otimes \chi^{c_1}{}_{c_2})\Delta$.

The $*$-structure on A induces a $*$-operation on the q-generators as follows:

$$(\chi^{a_1}{}_{a_2})^*(a) = \overline{\chi^{a_2}{}_{a_1}(\kappa^{-1}(a^*))}, \quad \forall a \in A \tag{44}$$

where the overline denotes the usual complex conjugation. As an example, let us consider the q-generators $\chi^a{}_b$ in the fundamental representation:

$$(\chi^{a_1}{}_{a_2})^{b_1}{}_{b_2} \equiv \chi^{a_1}{}_{a_2}(T^{b_1}{}_{b_2}) \tag{45}$$

By using eqs. (15) and (44), it is easy to find that

$$[(\chi^{a_1}{}_{a_2})^*]^{b_1}{}_{b_2} = \overline{(\chi^{a_2}{}_{a_1})^{b_2}{}_{b_1}} \tag{46}$$

$$\Rightarrow (\chi^{a_1}{}_{a_2})^* = (\chi^{a_2}{}_{a_1})^\dagger \tag{47}$$

where the dagger † means usual hermitian conjugation of the matrix that represents $\chi^{a_2}{}_{a_1}$.

The $GL_q(3)$ Lie algebra relations are presented in Table 4, with the standard index convention already used for the $T^a{}_b$ and the $\omega_a{}^b$. In the limit $q \to 1$, we recover the $SU(3)\otimes U(1)$ Lie algebra in the Cartan basis. The two commuting $U(1)$'s inside $SU(3)$ are generated by $\frac{1}{2\sqrt{3}}(\chi_1 - \chi_5)$ and $\frac{1}{6}(\chi_1 + \chi_5 - 2\chi_9)$, and the remaining $U(1)$ by $\chi_1 + \chi_5 + \chi_9$. The generators $\frac{1}{\sqrt{6}}(\chi_2, \chi_3, \chi_6, \chi_4, \chi_7, \chi_8)$ correspond to the roots of the $SU(3)$ hexagon, ordered counterclockwise, with χ_2 on the positive x-axis. Note that the q-generators $\chi_5, \chi_9, \chi_6, \chi_8$ close on the $GL_q(2)$ q-Lie algebra discussed for ex. in ref. [21] (there the generators are denoted by $\chi_1, \chi_2, \chi_+, \chi_-$). In the limit $q \to 1$, the rule (47) can be verified to yield the familiar relations $\chi_2 = (\chi_4)^\dagger$ etc.

The Cartan-Maurer equations are found by applying to $\omega_{c_1}{}^{c_2}$ the exterior differential as defined in (30):

$$d\omega_{c_1}{}^{c_2} = \frac{1}{q-q^{-1}}(\omega_b{}^b \wedge \omega_{c_1}{}^{c_2} + \omega_{c_1}{}^{c_2} \wedge \omega_b{}^b) \equiv -C^{a_1}{}_{a_2}{}^{b_1}{}_{b_2}|_{c_1}^{c_2}\ \omega_{a_1}{}^{a_2} \wedge \omega_{b_1}{}^{b_2}. \tag{48}$$

In order to obtain an explicit expression for the C structure constants in (48) which reduces to the correct $q = 1$ limit, we must use the relation (18) for the commutations of $\omega_{a_1}{}^{a_2}$ with $\omega_{b_1}{}^{b_2}$. Then the term $\omega_{c_1}{}^{c_2} \wedge \omega_b{}^b$ in (48) can be written as $-Z\omega\omega$ via formula (18), and we find the C-structure constants to be:

$$\begin{aligned} C^{a_1}{}_{a_2}{}^{b_1}{}_{b_2}|_{c_1}^{c_2} &= -\frac{1}{q-q^{-1}}(\delta^{a_1}_{a_2}\delta^{b_1}_{c_1}\delta^{c_2}_{b_2} - \frac{1}{q^2+q^{-2}}[\Lambda_{c_1}{}^{c_2}{}_b{}^b|^{a_1}{}_{a_2}{}^{b_1}{}_{b_2} + (\Lambda^{-1})_{c_1}{}^{c_2}{}_b{}^b|^{a_1}{}_{a_2}{}^{b_1}{}_{b_2}]) \\ &= -\frac{1}{q-q^{-1}}(\delta^{a_1}_{a_2}\delta^{b_1}_{c_1}\delta^{c_2}_{b_2} - \frac{1}{q^2+q^{-2}}[\delta^{a_1}_{a_2}\delta^{b_1}_{c_1}\delta^{c_2}_{b_2} + (\Lambda^{-1})_{c_1}{}^{c_2}{}_b{}^b|^{a_1}{}_{a_2}{}^{b_1}{}_{b_2}]), \end{aligned} \tag{49}$$

where we have used (see for example [21])

$$\Lambda_{c_1}{}^{c_2}{}_b{}^b\big|^{a_1}{}_{a_2}{}^{b_1}{}_{b_2} = \delta^{a_1}_{a_2}\delta^{b_1}_{c_1}\delta^{c_2}_{b_2}. \tag{50}$$

By considering the analogue of (17) for Λ^{-1}, it is not difficult to see that the terms proportional to $\frac{1}{q-q^{-1}}$ cancel, and the $q \to 1$ limit of (49) is well defined.

The Cartan-Maurer equations for $GL_q(3)$ are collected in Table 5.

Finally, we compute the Lie derivative of the q-group elements T_i and of the left-invariant 1-forms ω^i. As discussed in [21], the functionals χ_i are the q-analogue of tangent vectors at the origin of the group, while $t_i = \chi * = (id \otimes \chi_i)\Delta$ are the q-analogue of left- invariant vector fields. The Lie derivative along $t = \chi *$ of a generic element $\tau \in \Gamma^{\otimes n}$ is defined by [21]

$$\ell_t(\tau) \equiv \chi * \tau \equiv (id \otimes \chi)\Delta_R(\tau) \tag{51}$$

We refer to [21] for a discussion of the properties of the quantum Lie derivative. The Lie derivative of ω^i is essential for defining the q-gauge variations of the q-potentials ω^i, see refs. [7] and [21]. Table 6 and Table 7 contain the exterior derivative of the $T^a{}_b$ elements of $GL_q(3)$, and the Lie derivative of the left-invariant $\omega_a{}^b$. The relevant formulas are (40) and:

$$\chi^{a_1}{}_{a_2} * T^{b_1}{}_{b_2} = \frac{1}{q-q^{-1}} T^{b_1}{}_d[s\,(R^{-1})^{da_1}{}_{fe}(R^{-1})^{ef}{}_{a_2b_2} - \delta^{a_1}_{a_2}\delta^d_{b_2}] \tag{52}$$

$$\chi^{a_1}{}_{a_2} * \omega_{b_1}{}^{b_2} = \omega_{c_1}{}^{c_2}\mathbf{C}^{c_1}{}_{c_2}{}^{a_1}{}_{a_2}\big|^{b_2}_{b_1} \tag{53}$$

cf. ref. [21]. In Table 7 we actually give the q-analogue of the gauge variations of the potentials ω^i, i.e. $\delta\omega^i \equiv \varepsilon^k \ell_{t_k}\omega^i$, where ε^k is a “q-gauge parameter”.
As in the classical case, the q-Lie derivative commutes with the exterior differental:

$$d\,\ell_t = \ell_t d \tag{54}$$

and is a representation of the q-Lie algebra, since [21]:

$$[\ell_{t_i}, \ell_{t_j}]_q = \ell_{[t_i,t_j]_q} \tag{55}$$

with $[t_i, t_j]_q \equiv [\chi_i, \chi_j]_q\, *$. The q-commutator $[\ ,\ ]_q$ is defined by the left-hand side of eq. (42).

We conclude by observing that the $GL_q(2)$ bicovariant differential calculus of ref. [21] can be obtained by “projecting” the $GL_q(3)$ formulas obtained in this Letter into their $GL_q(2)$ subspace. This is done by rescaling the T, ω and χ with indices $1, 2, 3, 4, 7$ and letting the rescaling parameter go to zero.

Table 1

The R matrix for $GL_q(3)$

$$R^{ab}{}_{cd} = \begin{pmatrix} q & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & \lambda & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & q & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & \lambda & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & \lambda & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & q \end{pmatrix}$$

where $\lambda = q - q^{-1}$.

Table 2

The commutation relations between the left-invariant ω^i

$$\omega^1\omega^1 = \omega^2\omega^2 = \omega^3\omega^3 = \omega^4\omega^4 = \omega^6\omega^6 = \omega^7\omega^7 = \omega^8\omega^8 = 0$$

$$\omega^1\omega^2 + \omega^2\omega^1 = 0$$

$$\omega^1\omega^3 + \omega^3\omega^1 = 0$$

$$\omega^1\omega^4 + \omega^4\omega^1 = 0$$

$$\omega^1\,\omega^5 + \omega^5\,\omega^1 + \frac{1}{2}(1-q^2)(\omega^4\,\omega^2 - \omega^2\,\omega^4) = 0$$

$$\omega^1\,\omega^6 + \omega^6\,\omega^1 + \frac{1}{2}(1-q^2)(\omega^4\,\omega^3 - q^{-1}\,\omega^3\,\omega^4) = 0$$

$$\omega^1\omega^7 + \omega^7\omega^1 = 0$$

$$\omega^1\,\omega^8 + \omega^8\,\omega^1 + \frac{1}{2}(1-q^2)(\omega^7\,\omega^2 - q^{-1}\,\omega^2\,\omega^7) = 0$$

$$\omega^1\,\omega^9 + \omega^9\,\omega^1 + \frac{1}{2}(1-q^2)(\omega^7\,\omega^3 - \omega^3\,\omega^7) + \frac{1}{2}(1-q^2)^2\,\omega^2\,\omega^4 = 0$$

$$\omega^2\,\omega^3 + q\,\omega^3\,\omega^2 = 0$$

$$(1+q^4)\,\omega^2\,\omega^4 + 2\,q^2\,\omega^4\,\omega^2 + (q^2-1)\,(\omega^1\,\omega^5 + \omega^5\,\omega^1) = 0$$

$$q^2\,\omega^2\,\omega^5 + \omega^5\,\omega^2 + (1-q^2)^2\,\omega^1\,\omega^2 + (1-q^2)\,\omega^2\,\omega^1 = 0$$

$$\omega^2\,\omega^6 + q^{-1}\,\omega^6\,\omega^2 + \frac{1}{2}(q^{-2}-1)\,(\omega^5\,\omega^3 - \omega^3\,\omega^5) + (q^2-1)\omega^1\,\omega^3 = 0$$

$$(1+q^4)\,\omega^2\,\omega^7 + 2\,q^3\,\omega^7\,\omega^2 + q(q^2-1)(\omega^1\,\omega^8 + \omega^8\,\omega^1) = 0$$

$$q\,\omega^2\,\omega^8 + \omega^2\,\omega^8 = 0$$

$$\omega^2\,\omega^9 + \omega^9\,\omega^2 + \frac{1}{2}(q-q^{-1})(q\,\omega^3\,\omega^8 - \omega^8\,\omega^3) + \frac{1}{2}q^2(1-q^2)^2\,\omega^1\,\omega^2 + \frac{1}{2}(1-q^2)^2\,\omega^2\,\omega^5 = 0$$

$$(1+q^4)\,\omega^3\,\omega^4 + 2\,q^3\,\omega^4\,\omega^3 + q(q^2-1)(\omega^1\,\omega^6 + \omega^6\,\omega^1) = 0$$

$$(1+q^4)\,\omega^3\,\omega^5 + 2\,q^2\,\omega^5\,\omega^3 + q(q^2-1)(\omega^2\,\omega^6 + q^{-1}\,\omega^6\,\omega^2) + q^2(1-q^2)^2\,\omega^1\omega^3 = 0$$

$$\omega^3\omega^6 + q^{-1}\,\omega^6\omega^3 = 0$$

$$(1+q^4)\,\omega^3\omega^7 + 2q^2\,\omega^7\omega^3 - (1-q^2)^2\,(\omega^1\omega^5+\omega^5\omega^1) + (q^2-1)(\omega^1\omega^9+\omega^9\omega^1) + 2q^2(1-q^2)\omega^4\omega^2 = 0$$

$$(1+q^4)\,\omega^3\omega^8 + 2q\,\omega^8\omega^3 - (1-q^2)^2\,\omega^2\omega^5 + (1-q^4)\,\omega^5\omega^2 + (q^2-1)(\omega^2\omega^9+\omega^9\omega^2) - 2q^2(1-q^2)\omega^1\omega^2 = 0$$

$$q^2(1+q^2)(\omega^3\omega^9+q^{-2}\,\omega^9\omega^3) - (1-q^2)(2-q^2+q^4)\,\omega^1\omega^3 + 2q(1-q^2)\omega^6\omega^2 + q^2(1-q^2)^2\omega^2\omega^6 = 0$$

$$(1+q^2)(\omega^4\omega^5 + q^2\,\omega^5\omega^4) + q^2(q^4-1)\,\omega^4\omega^1 = 0$$

$$\omega^4\omega^6 + q\,\omega^6\omega^4 = 0$$

$$\omega^4\omega^7 + q^{-1}\,\omega^7\omega^4 = 0$$

$$\omega^4\omega^8 + q\omega^8\omega^4 + \frac{1}{2}q^{-1}(q^2-1)(\omega^5\omega^7 - \omega^7\omega^5) + q(q^2-1)\omega^7\omega^1 = 0$$

$$\omega^4\omega^9 + q\omega^9\omega^4 + \frac{1}{2}q^{-1}(q^2-1)(\omega^6\omega^7 - q^{-1}\,\omega^7\omega^6) = 0$$

$$(1+q^2)^2\,\omega^5\omega^5 + (1-q^2)^2\,(\omega^1\omega^5 + \omega^5\omega^1) + 2q^2(1-q^2)(\omega^2\omega^4 - \omega^4\omega^2) = 0$$

$$(1+q^2)(\omega^5\omega^6+\omega^6\omega^5) + (1-q^2)^2(\omega^1\omega^6+\omega^6\omega^1) + 2q(1-q^2)\omega^3\omega^4 - q^2(q^2-1)^2\omega^4\omega^3 = 0$$

$$(1+q^4)\,\omega^5\omega^7 + 2q^2\,\omega^7\omega^5 + q^2(1-q^2)^2\omega^7\omega^1 + q^2(q^2-1)(q^{-1}\,\omega^4\omega^8 + \omega^8\omega^4) = 0$$

$$\omega^5\omega^8 + \omega^8\omega^5 + (q^2-1)\,\omega^7\omega^2 = 0$$

$$\omega^5\omega^9+\omega^9\omega^5+\frac{1}{2}q^2(q^2-1)^2\,\omega^4\omega^2-\frac{1}{2}(q^2-1)^2\,\omega^7\omega^3+\frac{1}{2}(1-q^2)^2\omega^5\omega^5+\frac{1}{2}(q^2-1)(\omega^6\omega^8-\omega^8\omega^6) = 0$$

$$(1+q^4)\omega^6\omega^7+2q\omega^7\omega^6+q^2(1-q^2)\,\omega^1\omega^4-(1-q^2)^2\omega^5\omega^4+(1-q^4)\,\omega^4\omega^5+(q^2-1)(\omega^4\omega^9+\omega^9\omega^4) = 0$$

$$(1+q^4)\,\omega^6\omega^8+2q^2\omega^8\omega^6+(1-q^2)^2(\omega^1\omega^5+\omega^5\omega^1)-2q^2(1-q^2)^2\,\omega^4\omega^2+2q^2(1-q^2)\omega^2\omega^4-$$

$$-2q^2(1-q^2)\,\omega^7\omega^3 + 2q^2(1-q^2)\omega^5\omega^5 + (q^2-1)(\omega^5\omega^9 + \omega^9\omega^5) = 0$$

$$q^2(q^2+1)(\omega^6\omega^9+q^{-2}\,\omega^9\omega^6)+(q^2-1)^2\,(\omega^1\omega^6+\omega^6\omega^1)+2q(1-q^2)\omega^3\omega^4+q^4(1-q^2)^2\,\omega^4\omega^3+$$

$$+(1-q^2-q^4+q^6)\omega^5\omega^6 + (1-q^4)\,\omega^6\omega^5 = 0$$

$$\omega^7\omega^8 + q\,\omega^8\omega^7 = 0$$

$$(1+q^2)(\omega^7\omega^9 + q^2\,\omega^9\omega^7) + q^4(q^4-1)\omega^7\omega^1 + q^2(q^4-1)\omega^8\omega^4 = 0$$

$$(1+q^2)(\omega^8\omega^9 + q^2\,\omega^9\omega^8) + q^4(q^4-1)\omega^7\omega^2 + q^2(q^4-1)\omega^8\omega^5 = 0$$

$$(1+q^2)^2\,\omega^9\omega^9+(1-q^2)^2(\omega^1\omega^9+\omega^9\omega^1)-2q^2(1-q^2)^2(\omega^2\omega^4+q^2\,\omega^4\omega^2)+2q^2(1-q^2)(\omega^3\omega^7-q^2\,\omega^7\omega^3)$$

$$-2q^2(1-q^2)^2\,\omega^5\omega^5 + (1-q^2)^2\,(\omega^5\omega^9 + \omega^9\omega^5) + 2q^2(q^2-1)(\omega^8\omega^6 - \omega^6\omega^8) = 0$$

Table 3

The commutation relations between ω^i and T_j

$$
\begin{array}{llll}
\omega^1T_1 = q^{-2}T_1\omega^1 & \omega^1T_2 = T_2\omega^1 & \omega^1T_3 = T_3\omega^1 & \omega^1T_4 = q^{-2}T_4\omega^1 \\
\omega^1T_5 = T_5\omega^1 & \omega^1T_6 = T_6\omega^1 & \omega^1T_7 = q^{-2}T_7\omega^1 & \omega^1T_8 = T_8\omega^1 \\
\omega^1T_9 = T_9\omega^1 & \omega^2T_1 = q^{-1}T_1\omega^2 & \omega^2T_3 = T_3\omega^2 & \omega^2T_4 = q^{-1}T_4\omega^2 \\
\omega^2T_6 = T_6\omega^2 & \omega^2T_7 = q^{-1}T_7\omega^2 & \omega^2T_9 = T_9\omega^2 & \omega^3T_1 = q^{-1}T_1\omega^3 \\
\omega^3T_2 = T_2\omega^3 & \omega^3T_4 = q^{-1}T_4\omega^3 & \omega^3T_5 = T_5\omega^3 & \omega^3T_7 = q^{-1}T_7\omega^3 \\
\omega^3T_8 = T_8\omega^3 & \omega^4T_2 = q^{-1}T_2\omega^4 & \omega^4T_3 = T_3\omega^4 & \omega^4T_5 = q^{-1}T_5\omega^4 \\
\omega^4T_6 = T_6\omega^4 & \omega^4T_8 = q^{-1}T_8\omega^4 & \omega^4T_9 = T_9\omega^4 & \omega^5T_3 = T_3\omega^5 \\
\omega^5T_6 = T_6\omega^5 & \omega^5T_9 = T_9\omega^5 & \omega^6T_2 = q^{-1}T_2\omega^6 & \omega^6T_5 = q^{-1}T_5\omega^6 \\
\omega^6T_8 = q^{-1}T_8\omega^6 & \omega^7T_3 = q^{-1}T_3\omega^7 & \omega^7T_6 = q^{-1}T_7\omega^6 & \omega^7T_9 = q^{-1}T_9\omega^7 \\
\omega^8T_3 = q^{-1}T_3\omega^8 & \omega^8T_6 = q^{-1}T_6\omega^8 & \omega^8T_9 = q^{-1}T_9\omega^8 &
\end{array}
$$

$$
\begin{array}{ll}
\omega^2T_2 = (q^{-2}-1)T_1\omega^1 + q^{-1}T_2\omega^2 & \omega^2T_5 = (q^{-2}-1)T_4\omega^1 + q^{-1}T_5\omega^2 \\
\omega^2T_8 = (q^{-2}-1)T_7\omega^1 + q^{-1}T_8\omega^2 & \omega^4T_1 = (q^{-2}-1)T_2\omega^1 + q^{-1}T_1\omega^4 \\
\omega^4T_4 = (q^{-2}-1)T_5\omega^1 + q^{-1}T_4\omega^4 & \omega^4T_7 = (q^{-2}-1)T_8\omega^1 + q^{-1}T_7\omega^4 \\
\omega^5T_1 = (q^{-1}-q)T_2\omega^2 + T_1\omega^5 & \omega^5T_4 = (q^{-1}-q)T_5\omega^2 + T_4\omega^5 \\
\omega^5T_7 = (q^{-1}-q)T_8\omega^2 + T_7\omega^5 & \omega^6T_1 = (q^{-1}-q)T_2\omega^3 + T_1\omega^6 \\
\omega^6T_4 = (q^{-1}-q)T_5\omega^3 + T_4\omega^6 & \omega^6T_7 = (q^{-1}-q)T_8\omega^3 + T_7\omega^6 \\
\omega^7T_1 = (q^{-2}-1)T_3\omega^1 + q^{-1}T_1\omega^7 & \omega^7T_2 = (q^{-1}-q)T_3\omega^4 + T_2\omega^7 \\
\omega^7T_4 = (q^{-2}-1)T_6\omega^1 + q^{-1}T_4\omega^7 & \omega^7T_5 = (q^{-1}-q)T_6\omega^4 + T_5\omega^7 \\
\omega^7T_7 = (q^{-2}-1)T_9\omega^1 + q^{-1}T_7\omega^7 & \omega^7T_8 = (q^{-1}-q)T_9\omega^4 + T_8\omega^7 \\
\omega^8T_4 = (q^{-1}-q)T_6\omega^2 + T_4\omega^8 & \omega^8T_7 = (q^{-1}-q)T_9\omega^2 + T_7\omega^8 \\
\omega^9T_1 = (q^{-1}-q)T_3\omega^3 + T_1\omega^9 & \omega^9T_2 = (q^{-1}-q)T_3\omega^6 + T_2\omega^9 \\
\omega^9T_4 = (q^{-1}-q)T_6\omega^3 + T_4\omega^9 & \omega^9T_5 = (q^{-1}-q)T_6\omega^6 + T_5\omega^9 \\
\omega^9T_7 = (q^{-1}-q)T_9\omega^3 + T_7\omega^9 & \omega^9T_8 = (q^{-1}-q)T_9\omega^6 + T_8\omega^9
\end{array}
$$

$$
\begin{array}{l}
\omega^3T_3 = (q^{-2}-1)T_1\omega^1 + (q^{-1}-q)T_2\omega^2 + q^{-1}T_3\omega^3 \\
\omega^3T_6 = (q^{-2}-1)T_4\omega^1 + (q^{-1}-q)T_5\omega^2 + q^{-1}T_6\omega^3 \\
\omega^3T_9 = (q^{-2}-1)T_7\omega^1 + (q^{-1}-q)T_8\omega^2 + q^{-1}T_9\omega^3 \\
\omega^5T_2 = (q^{-1}-q)^2\,T_2\omega^1 + (q^{-1}-q)T_1\omega^4 + q^{-2}T_2\omega^5 \\
\omega^5T_5 = (q^{-1}-q)^2\,T_5\omega^1 + (q^{-1}-q)T_4\omega^4 + q^{-2}T_5\omega^5 \\
\omega^5T_8 = (q^{-1}-q)^2\,T_8\omega^1 + (q^{-1}-q)T_7\omega^4 + q^{-2}T_8\omega^5 \\
\omega^6T_3 = (q^{-1}-q)^2T_2\omega^1 + (q^{-1}-q)T_1\omega^4 + (q^{-2}-1)T_2\omega^5 + q^{-1}T_3\omega^6 \\
\omega^6T_6 = (q^{-1}-q)^2T_5\omega^1 + (q^{-1}-q)T_4\omega^4 + (q^{-2}-1)T_5\omega^5 + q^{-1}T_6\omega^6 \\
\omega^6T_9 = (q^{-1}-q)^2T_8\omega^1 + (q^{-1}-q)T_7\omega^4 + (q^{-2}-1)T_8\omega^5 + q^{-1}T_9\omega^6 \\
\omega^8T_2 = (q^{-1}-q)^2T_3\omega^1 + (q^{-1}-q)T_1\omega^7 + (q^{-2}-1)T_3\omega^5 + q^{-1}T_2\omega^8 \\
\omega^8T_5 = (q^{-1}-q)^2T_6\omega^1 + (q^{-1}-q)T_4\omega^7 + (q^{-2}-1)T_6\omega^5 + q^{-1}T_5\omega^8 \\
\omega^8T_8 = (q^{-1}-q)^2T_9\omega^1 + (q^{-1}-q)T_7\omega^7 + (q^{-2}-1)T_9\omega^5 + q^{-1}T_8\omega^8 \\
\omega^9T_3 = (q^{-1}-q)^2T_3\omega^1 + (q^{-1}-q)^2T_3\omega^5 + (q^{-1}-q)T_1\omega^7 + (q^{-1}-q)T_2\omega^8 + q^{-2}T_3\omega^9 \\
\omega^9T_6 = (q^{-1}-q)^2T_6\omega^1 + (q^{-1}-q)^2T_6\omega^5 + (q^{-1}-q)T_4\omega^7 + (q^{-1}-q)T_5\omega^8 + q^{-2}T_6\omega^9 \\
\omega^9T_9 = (q^{-1}-q)^2T_9\omega^1 + (q^{-1}-q)^2T_9\omega^5 + (q^{-1}-q)T_7\omega^7 + (q^{-1}-q)T_8\omega^8 + q^{-2}T_9\omega^9
\end{array}
$$

Table 4

The q-Lie algebra

$$
\begin{aligned}
&\chi_1\chi_2 - \chi_2\chi_1 + (1-q^2)(\chi_2\chi_5 + \chi_3\chi_8) = q\chi_2 \\
&\chi_1\chi_3 - \chi_3\chi_1 + (1-q^2)(q^2\chi_9\chi_3 + \chi_2\chi_6) = q^3\chi_3 \\
&\chi_1\chi_4 - \chi_4\chi_1 - (1-q^2)(\chi_5\chi_4 + \chi_6\chi_7) = -q\chi_4 \\
&\chi_1\chi_5 - \chi_5\chi_1 = 0 \\
&\chi_1\chi_6 - \chi_6\chi_1 = 0 \\
&\chi_1\chi_7 - \chi_7\chi_1 - (1-q^2)(\chi_8\chi_4 + \chi_9\chi_7) = -q\chi_7 \\
&\chi_1\chi_8 - \chi_8\chi_1 = 0 \\
&\chi_1\chi_9 - \chi_9\chi_1 = 0 \\
&\chi_2\chi_3 - q^{-1}\chi_3\chi_2 = 0 \\
&\chi_2\chi_4 - \chi_4\chi_2 + (1-q^2)[\chi_5\chi_1 - \chi_5\chi_5 + \chi_7\chi_3 - q^2\chi_8\chi_6 - (1-q^2)(\chi_9\chi_1 - \chi_9\chi_5)] \\
&\hspace{20em} = q^3(\chi_1 - \chi_5) \\
&\chi_2\chi_5 - q^2\chi_5\chi_2 + (1-q^2)\chi_3\chi_8 = q\chi_2 \\
&\chi_2\chi_6 - q\chi_6\chi_2 + (1-q^2)\chi_3\chi_9 = q\chi_3 \\
&\chi_2\chi_7 - q^{-1}\chi_7\chi_2 + (q^{-1}-q)(\chi_8\chi_1 - \chi_8\chi_5 - \chi_9\chi_8) = -\chi_8 \\
&\chi_2\chi_8 - q\chi_8\chi_2 = 0 \\
&\chi_2\chi_9 - \chi_9\chi_2 = 0 \\
&\chi_3\chi_4 - q^{-1}\chi_4\chi_3 + (q^{-1}-q)(\chi_6\chi_1 - \chi_5\chi_6 - \chi_6\chi_9) = -\chi_6 \\
&\chi_3\chi_5 - \chi_5\chi_3 + (q^{-1}-q)\chi_6\chi_2 = 0 \\
&\chi_3\chi_6 - q\chi_6\chi_3 = 0 \\
&\chi_3\chi_7 - \chi_7\chi_3 - (1-q^2)(\chi_8\chi_6 - \chi_9\chi_1 + \chi_9\chi_9) = q(\chi_1 - \chi_9) \\
&\chi_3\chi_8 - q\chi_8\chi_3 + (1-q^2)\chi_9\chi_2 = q\chi_2 \\
&\chi_3\chi_9 - q^2\chi_9\chi_3 = q\chi_3 \\
&\chi_4\chi_5 - q^{-2}\chi_5\chi_4 + (1-q^{-2})\chi_6\chi_7 = -q^{-1}\chi_4 \\
&\chi_4\chi_6 - q^{-1}\chi_6\chi_4 = 0 \\
&\chi_4\chi_7 - q\chi_7\chi_4 = 0 \\
&\chi_4\chi_8 - q^{-1}\chi_8\chi_4 - (q^{-1}-q)\chi_9\chi_7 = -\chi_7 \\
&\chi_4\chi_9 - \chi_9\chi_4 = 0 \\
&\chi_5\chi_6 - \chi_6\chi_5 + (1-q^2)\chi_6\chi_9 = q\chi_6 \\
&\chi_5\chi_7 - \chi_7\chi_5 + (q^{-1}-q)\chi_4\chi_8 = 0 \\
&\chi_5\chi_8 - \chi_8\chi_5 - (1-q^2)\chi_9\chi_8 = -q\chi_8 \\
&\chi_5\chi_9 - \chi_9\chi_5 = 0 \\
&\chi_6\chi_7 - q\chi_7\chi_6 + (1-q^2)\chi_9\chi_4 = q\chi_4 \\
&\chi_6\chi_8 - \chi_8\chi_6 + (1-q^2)(\chi_9\chi_5 - \chi_9\chi_9) = q(\chi_5 - \chi_9) \\
&\chi_6\chi_9 - q^2\chi_9\chi_6 = q\chi_6 \\
&\chi_7\chi_8 - q^{-1}\chi_8\chi_7 = 0 \\
&\chi_7\chi_9 - q^{-2}\chi_9\chi_7 = q^{-1}\chi_7 \\
&\chi_8\chi_9 - q^{-2}\chi_9\chi_8 = -q^{-1}\chi_8
\end{aligned}
$$

Table 5

The Cartan-Maurer equations

$$\begin{aligned}
(1+q^4)\,d\omega^1 = {} & -q\,(1-q^2)\,\omega^1\omega^5 - q\,(1-q^2)\,\omega^1\omega^9 - q^5\omega^2\omega^4 - q^3\omega^3\omega^7 \\
& +q^3\omega^4\omega^2 - q(1-q^2)\,\omega^5\omega^1 + q^3\,\omega^7\omega^3 - q\,(1-q^2)\,\omega^9\omega^1 \\
(1+q^4)\,d\omega^2 = {} & -q^7\,\omega^1\omega^2 + q^3\,\omega^2\omega^1 - q\,(1-q^2+q^4)\,\omega^2\omega^5 - q\,(1-q^2)\,\omega^2\omega^9 \\
& -q^3\,\omega^3\omega^8 + q^3\,\omega^5\omega^2 + q^2\,\omega^8\omega^3 - q\,(1-q^2)\,\omega^9\omega^2 \\
(1+q^4)\,d\omega^3 = {} & -q^7\,\omega^1\omega^3 - q^5\,\omega^2\omega^6 + q^3\,\omega^3\omega^1 - q\,(1-q^2)\,\omega^3\omega^5 - q\,\omega^3\omega^9 + q^2\,\omega^6\omega^2 + q^3\,\omega^9\omega^3 \\
(1+q^4)\,d\omega^4 = {} & q^3\,\omega^1\omega^4 - q^7\,\omega^4\omega^1 + q^3\,\omega^4\omega^5 - q\,(1-q^2)\,\omega^4\omega^9 \\
& -q\,(1-q^2+q^4)\,\omega^5\omega^4 - q^3\,\omega^6\omega^7 + q^2\,\omega^7\omega^6 - q\,(1-q^2)\,\omega^9\omega^4 \\
(1+q^4)\,d\omega^5 = {} & q^3\,\omega^2\omega^4 - q^3\,(1-q^2+q^4)\,\omega^4\omega^2 - q\,(-1+q^2)^2\,\omega^5\omega^5 - q\,(1-q^2)\,\omega^5\omega^9 \\
& -q^3\,\omega^6\omega^8 - q^3\,(1-q^2)\,\omega^7\omega^3 + q^3\,\omega^8\omega^6 - q\,(1-q^2)\,\omega^9\omega^5 \\
(1+q^4)\,d\omega^6 = {} & q^2\,\omega^3\omega^4 - q^7\,\omega^4\omega^3 - q^5\,\omega^5\omega^6 + q^3\,\omega^6\omega^5 - q\,\omega^6\omega^9 + q^3\,\omega^9\omega^6 \\
(1+q^4)\,d\omega^7 = {} & q^3\,\omega^1\omega^7 + q^2\,\omega^4\omega^8 - q\,(1-q^2)\,\omega^5\omega^7 - q^7\,\omega^7\omega^1 \\
& +q^3\,\omega^7\omega^9 - q^5\,\omega^8\omega^4 - q\,\omega^9\omega^7 \\
(1+q^4)\,d\omega^8 = {} & q^2\,\omega^2\omega^7 + q^3\,\omega^5\omega^8 - q^7\,\omega^7\omega^2 - q^5\,\omega^8\omega^5 + q^3\,\omega^8\omega^9 - q\,\omega^9\omega^8 \\
(1+q^4)\,d\omega^9 = {} & -q^3\,(1-q^2)\,\omega^2\omega^4 + q^3\,\omega^3\omega^7 - q^5\,(1-q^2)\,\omega^4\omega^2 - q^3\,(1-q^2)\,\omega^5\omega^5 \\
& +q^3\,\omega^6\omega^8 - q^5\,\omega^7\omega^3 - q^3\,\omega^8\omega^6 - q\,(1-q^2)\,\omega^9\omega^9
\end{aligned}$$

Table 6

The exterior derivative of T_i

$$d\,T_1 = \frac{(-q^2+s)}{-q+q^3}\,T_1\,\omega^1 - s\,T_2\,\omega^2 - s\,T_3\,\omega^3 + \frac{(-1+s)}{-q^{-1}+q}\,T_1\,(\omega^5+\omega^9)$$

$$d\,T_2 = \frac{(-q^2+s-q^2\,s+q^4\,s)}{-q+q^3}\,T_2\,\omega^1 - s\,T_1\,\omega^4 + \frac{(-q^2+s)}{-q+q^3}\,T_2\,\omega^5 - s\,T_3\,\omega^6 + \frac{(-1+s)}{-q^{-1}+q}\,T_2\,\omega^9$$

$$d\,T_3 = \frac{(-q^2+s-q^2\,s+q^4\,s)}{-q+q^3}\,T_3\,(\omega^1+\omega^5) - s\,T_1\,\omega^7 - s\,T_2\,\omega^8 + \frac{(-q^2+s)}{-q+q^3}\,T_3\,\omega^9$$

$$d\,T_4 = \frac{(-q^2+s)}{-q+q^3}\,T_4\,\omega^1 - s\,T_5\,\omega^2 - s\,T_6\,\omega^3 + \frac{(-1+s)}{-q^{-1}+q}\,T_4\,(\omega^5+\omega^9)$$

$$d\,T_5 = \frac{(-q^2+s-q^2\,s+q^4\,s)}{-q+q^3}\,T_5\,\omega^1 - s\,T_4\,\omega^4 + \frac{(-q^2+s)}{-q+q^3}\,T_5\,\omega^5 - s\,T_6\,\omega^6 + \frac{(-1+s)}{-q^{-1}+q}\,T_5\,\omega^9$$

$$d\,T_6 = \frac{(-q^2+s-q^2\,s+q^4\,s)}{-q+q^3}\,T_6\,(\omega^1+\omega^5) - s\,T_4\,\omega^7 - s\,T_5\,\omega^8 + \frac{(-q^2+s)}{-q+q^3}\,T_6\,\omega^9$$

$$d\,T_7 = \frac{(-q^2+s)}{-q+q^3}\,T_7\,\omega^1 - s\,T_8\,\omega^2 - s\,T_9\,\omega^3 + \frac{(-1+s)}{-q^{-1}+q}\,T_7\,(\omega^5+\omega^9)$$

$$d\,T_8 = \frac{(-q^2+s-q^2\,s+q^4\,s)}{-q+q^3}\,T_8\,\omega^1 - s\,T_7\,\omega^4 + \frac{(-q^2+s)}{-q+q^3}\,T_8\,\omega^5 - s\,T_9\,\omega^6 + \frac{(-1+s)}{-q^{-1}+q}\,T_8\,\omega^9$$

$$d\,T_9 = \frac{(-q^2+s-q^2\,s+q^4\,s)}{-q+q^3}\,T_9\,(\omega^1+\omega^5) - s\,T_7\,\omega^7 - s\,T_8\,\omega^8 + \frac{(-q^2+s)}{-q+q^3}\,T_9\,\omega^9$$

Table 7

The Lie derivative of ω^i

$$\epsilon^i\,\ell_{t_i}\,\omega^1 = q^3\,\epsilon^4\,\omega^2 + q\,\epsilon^7\,\omega^3 - q\,\epsilon^2\,\omega^4 - q\,\epsilon^3\,\omega^7 + \epsilon^1\left[(-q+q^5)\,\omega^1 + (q^{-1}-q)\,\omega^5 + (q^{-1}-q)\,\omega^9\right]$$

$$\epsilon^i\,\ell_{t_i}\,\omega^2 = -q^{-1}\,\epsilon^1\,\omega^2 + q^3\,\epsilon^5\,\omega^2 + q\,\epsilon^8\,\omega^3 - \epsilon^3\,\omega^8 + \epsilon^2\left[q^5\,\omega^1 - q\,\omega^5 + (q^{-1}-q)\,\omega^9\right]$$

$$\epsilon^i\,\ell_{t_i}\,\omega^3 = q^3\,\epsilon^6\,\omega^2 - q^{-1}\,\epsilon^1\,\omega^3 + q\,\epsilon^9\,\omega^3 - \epsilon^2\,\omega^6 + \epsilon^3\,(q^5\,\omega^1 - q\,\omega^9)$$

$$\epsilon^i\,\ell_{t_i}\,\omega^4 = (-q^{-1}+q+q^5)\,\epsilon^1\,\omega^4 - q^{-1}\,\epsilon^5\,\omega^4 + q\,\epsilon^7\,\omega^6 - \epsilon^6\,\omega^7 + \epsilon^4\left[-q\,\omega^1 + (q^{-1}-q+q^3)\,\omega^5 + (q^{-1}-q)\,\omega^9\right]$$

$$\begin{aligned}\epsilon^i\,\ell_{t_i}\,\omega^5 = {}& -q\,\epsilon^4\,\omega^2 + (q-q^3+q^5)\,\epsilon^2\,\omega^4 + \epsilon^1\left((q-q^3)\,\omega^1 + (-q^{-1}+q)\,\omega^5\right) + q\,\epsilon^8\,\omega^6 + \\ & (q-q^3)\,\epsilon^3\,\omega^7 - q\,\epsilon^6\,\omega^8 + \epsilon^5\left[(-q+q^3)\,\omega^5 + (q^{-1}-q)\,\omega^9\right]\end{aligned}$$

$$\epsilon^i\,\ell_{t_i}\,\omega^6 = -\epsilon^4\,\omega^3 + q^5\,\epsilon^3\,\omega^4 + (-q^{-1}+q)\,\epsilon^1\,\omega^6 - q^{-1}\,\epsilon^5\,\omega^6 + q\,\epsilon^9\,\omega^6 + \epsilon^6\,(q^3\,\omega^5 - q\,\omega^9)$$

$$\begin{aligned}\epsilon^i\,\ell_{t_i}\,\omega^7 = {}& -\epsilon^8\,\omega^4 + (-q^{-1}+q+q^5)\,\epsilon^1\,\omega^7 + (-q^{-1}+q)\,\epsilon^5\,\omega^7 - q^{-1}\,\epsilon^9\,\omega^7 + \\ & q^3\,\epsilon^4\,\omega^8 + \epsilon^7\left[-q\,\omega^1 + (q^{-1}-q)\,\omega^5 + q^{-1}\,\omega^9\right]\end{aligned}$$

$$\epsilon^i\,\ell_{t_i}\,\omega^8 = -\epsilon^7\,\omega^2 + q^5\,\epsilon^2\,\omega^7 + (-q^{-1}+q)\,\epsilon^1\,\omega^8 + (-q^{-1}+q+q^3)\,\epsilon^5\,\omega^8 - q^{-1}\,\epsilon^9\,\omega^8 + \epsilon^8\,(-q\,\omega^5 + q^{-1}\,\omega^9)$$

$$\begin{aligned}\epsilon^i\,\ell_{t_i}\,\omega^9 = {}& (q-q^3)\,\epsilon^4\,\omega^2 - q\,\epsilon^7\,\omega^3 + (q^3-q^5)\,\epsilon^2\,\omega^4 - q\,\epsilon^8\,\omega^6 + q^3\,\epsilon^3\,\omega^7 + q\,\epsilon^6\,\omega^8 + \\ & \epsilon^1\left[(q^3-q^5)\,\omega^1 + (-q^{-1}+q)\,\omega^9\right] + \epsilon^5\left[(q-q^3)\,\omega^5 + (-q^{-1}+q)\,\omega^9\right]\end{aligned}$$

References

[1] V. Drinfeld, Sov. Math. Dokl. **32** (1985) 254.

[2] M. Jimbo, Lett. Math. Phys. **10** (1985) 63; **11** (1986) 247.

[3] L.D. Faddeev, N.Yu. Reshetikhin and L.A. Takhtajan, Algebra and Analysis, **1** (1987) 178.

[4] S. Majid, Int. J. Mod. Phys. **A5** (1990) 1.

[5] A.P. Isaev and Z. Popowicz, *q-Trace for the Quantum Groups and q-deformed Yang-Mills Theory*, Wroclaw preprint ITP UWr 786-91.

[6] I.Y. Aref'eva and I.V. Volovich, Mod. Phys. Lett **A6** (1991) 893; Phys. Lett. **B264** (1991) 62.

[7] L. Castellani, *Gauge theories of quantum groups*, Torino preprint DFTT 19/92, April 1992, to be published in Phys. Lett. B.

[8] T. Brzeziński and S. Majid, *Quantum group gauge theory on quantum spaces*, preprint DAMPT/92-27.

[9] L. Castellani, Phys. Lett. **B279** (1992) 291.

[10] S.L. Woronowicz, Publ. RIMS, Kyoto Univ., Vol. **23**, 1 (1987) 117; Commun. Math. Phys. **111** (1987) 613 and Commun. Math. Phys. **122**, (1989) 125.

[11] Yu.I. Manin, *Quantum Groups and Non-Commutative Geometry*, Université de Montréal, Centre de Recherches Mathématiques, 1988.

[12] D. Bernard, *Quantum Lie algebras and differential calculus on quantum groups*, Proc. 1990 Yukawa Int. Seminar, Kyoto; Phys. Lett. **260B** (1991) 389.

[13] P. Podleś and S.L. Woronowicz, Commun. Math. Phys. **130** (1990) 381.

[14] B. Zumino, in *Recent Advances in Field Theories*, Annecy meeting in honour of R. Stora, 1990;

J. Wess and B. Zumino, Nucl. Phys. (Proc. Suppl.) **B18** (1990) 302.

W.M. Schmidke, S.P. Vokos and B. Zumino, Z. Phys. C,**48** (1990) 249.

A. Schirrmacher, J. Wess and B. Zumino, Z. Phys. C **49** (1991) 317.

[15] W. Weich, *Die Quantengruppe $SU_q(2)$ - Kovariante Differentialrechnung und ein quantenmechanisches Modell*, Ph.D. Thesis, Karlsruhe University (1990).

[16] U. Carow-Watamura, M. Schlieker and S. Watamura, Z. Phys. C, **49** (1991) 439.

U. Carow-Watamura, M. Schlieker, S. Watamura and W. Weich, Commun. Math. Phys. **142** (1991) 605.

[17] B. Jurčo, Lett. Math. Phys. **22** (1991) 177.

[18] B. Zumino, *Introduction to the Differential Geometry of Quantum Groups*, LBL-31432 and UCB-PTH-62/91, notes of a plenary talk given at the 10-th IAMP Conf., Leipzig (1991).

[19] F. Müller-Hoissen, J. Phys. A **25** (1992) 1703.

[20] X.D. Sun and S.K. Wang, *Bicovariant differential calculus on quantum group* $GL_q(n)$, Worldlab-Beijing preprint CCAST-92-04, ASIAM-92-07, ASITP-92-12 (1992).

[21] P. Aschieri and L. Castellani, *An introduction to non-commutative differential geometry on quantum groups*, preprint CERN-TH.6565/92, DFTT-22/92.

[22] P. Schupp, P. Watts and B. Zumino, *Differential Geometry on Linear Quantum Groups*, preprint LBL-32314, UCB-PTH-92/13.